

UV finiteness of 3D Yang-Mills theories with a regulating mass in the Landau gauge

D. Dudal^{a,*}, J.A. Gracey^{b,†}, R.F. Sobreiro^{c,‡}, S.P. Sorella^{c,§} and H. Verschelde^{a,¶}

^a *Ghent University
Department of Mathematical Physics and Astronomy
Krijgslaan 281-S9
B-9000 Gent, Belgium*

^b *Theoretical Physics Division
Department of Mathematical Sciences
University of Liverpool
P.O. Box 147, Liverpool, L69 3BX, United Kingdom*

^c *UERJ - Universidade do Estado do Rio de Janeiro
Rua São Francisco Xavier 524, 20550-013 Maracanã
Rio de Janeiro, Brasil*

We prove that three-dimensional Yang-Mills theories in the Landau gauge supplemented with a infrared regulating, parity preserving mass term are ultraviolet finite to all orders. We also extend this result to the Curci-Ferrari gauge.

PACS numbers:

I. INTRODUCTION

In a previous paper [1], we started to investigate three-dimensional Yang-Mills theories in the Landau gauge, supplemented with a parity preserving mass term. We recall that the $SU(N)$ Yang-Mills action in d -dimensional Euclidean space time is given by

$$S = S_{\text{YM}} + S_{\text{GF+FP}} , \quad (1)$$

where the classical action reads

$$\begin{aligned} S_{\text{YM}} &= \int d^d x \left(\frac{1}{4} F_{\mu\nu}^a F_{\mu\nu}^a \right) , \\ F_{\mu\nu}^a &= \partial_\mu A_\nu^a - \partial_\nu A_\mu^a - g f^{abc} A_\mu^b A_\nu^c , \end{aligned} \quad (2)$$

and the gauge fixing part

$$S_{\text{GF+FP}} = \int d^d x \left(b^a \partial_\mu A_\mu^a + c^a \partial_\mu D_\mu^{ab} \bar{c}^b \right) , \quad (3)$$

enforces the Landau gauge $\partial_\mu A_\mu^a = 0$. b^a is the Lagrange multiplier and $c^a, \bar{c}^a$ are the ghost and antighost fields. The covariant derivative in the adjoint representation is given by

$$D_\mu^{ab} = \delta^{ab} \partial_\mu - g f^{abc} A_\mu^c . \quad (4)$$

For $d = 3$, the coupling g^2 becomes a dimensional quantity itself. As a consequence, the three-dimensional Yang-Mills theory (1) is superrenormalizable. Such theories exhibit serious problems in the infrared sector, as extensively

*D. Dudal is a postdoctoral fellow of the *Special Research Fund* of Ghent University; Electronic address: david.dudal@ugent.be

†Electronic address: jag@amtp.liv.ac.uk

‡Electronic address: sobreiro@dft.if.uerj.br

§Work supported by FAPERJ, Fundação de Amparo à Pesquisa do Estado do Rio de Janeiro, under the program *Cientista do Nosso Estado*, E-26/151.947/2004.; Electronic address: sorella@uerj.br

¶Electronic address: henri.verschelde@ugent.be

discussed in [2]. The origin of this problem is easy to understand. If we were to write down a perturbative series in the massive coupling g^2 , then at a certain point inverse powers of the external momenta will inevitably appear in order to obtain a certain desired dimensionality. Consequently, at increasing order of perturbation theory, the low momentum region becomes more and more problematic to consider.

A natural solution to this problem might be the dynamical generation of a mass m . If this were the case, a perturbative expansion in the dimensionless parameter $\frac{g^2}{m}$ might emerge, ensuring a safe infrared limit. A few mechanisms and techniques behind such a mass generation are presented in for example [3, 4, 5, 6, 7, 8]

A common feature of the approaches of for example [3, 4, 5, 6] is that the necessary calculations are performed with a massive version of Yang-Mills in a certain gauge. Essentially, a nonperturbative mass is derived by constructing a gap equation, in a particular approximation scheme, with nontrivial solutions. This necessitates in fact a proof of the renormalizability of the massive Yang-Mills theories in the specific gauges used [25], as the employed gap equation should be at least renormalizable.

In [1], we have already shown that the Yang-Mills action (1) in the Landau gauge, supplemented with the parity preserving mass term $\frac{1}{2}m^2 A_\mu^2$

$$S = S_{\text{YM}} + S_{\text{GF+FP}} + \int d^d x \frac{1}{2} m^2 A_\mu^a A_\mu^a, \quad (5)$$

is renormalizable to all orders using the algebraic renormalization formalism [9]. This general proof was possible since the action (5) enjoys a certain number of Ward identities, even in the massive case, allowing one to sufficiently restrict the most general allowed counterterm, so that it can be reabsorbed by a multiplicative redefinition of the fields and parameters present in the original action [1]. When we define the bare fields and parameters by

$$\begin{aligned} A_{0\mu}^a &= Z_A^{1/2} A_\mu^a, & c_0^a &= Z_c^{1/2} c^a, \\ g_0 &= Z_g g, & m_0^2 &= Z_m^2 m^2, \\ \bar{c}_0^a &= Z_{\bar{c}}^{1/2} \bar{c}^a, & b_0^a &= Z_b^{1/2} b^a, \end{aligned} \quad (6)$$

then it turns out that only Z_A and Z_g are independent, while all the other renormalization factors can be expressed in terms of these by the following relations,

$$\begin{aligned} Z_{\bar{c}} &= Z_c = Z_g^{-1} Z_A^{-1/2}, \\ Z_b &= Z_A^{-1}, \\ Z_{m^2} &= Z_g Z_A^{-1/2}, \end{aligned} \quad (7)$$

which follow from the Ward identities [1]. We recall that the operator $\int d^3 x A_\mu^2$ is BRST-invariant on-shell [10].

This situation is completely similar to four dimensions [11]. We especially want to focus attention on the fact that the mass m does not renormalize independently.

Although the algebraic formalism enables us to prove the renormalizability to all orders, it does not allow us to say anything about the potential ultraviolet finiteness of the theory (5). In [1], we already pointed out that the theory is ultraviolet finite at the one loop level, when dimensional regularization is employed. In this paper, we extend this analysis to two loop order, by explicitly verifying the absence of ultraviolet divergences. Here, we want to stress that *individual* diagrams can be divergent. Nevertheless, when taken together to obtain the renormalization factors, it turns out that no divergent terms remain, and hence the theory is also completely finite at two loop order. Notice that it would be sufficient to consider the renormalization of for example the gluon and ghost two point function, as only two independent renormalization factors are needed.

By a power counting argument, we shall consequently deduce that the theory is ultraviolet finite to any order.

To our knowledge, a complete proof of the ultraviolet finiteness of three-dimensional Yang-Mills theories using a renormalizable gauge and including a regulating mass compatible with a Slavnov-Taylor identity [1], is still lacking so far.

Most works remained at the one loop level, not encountering any divergences in three dimensions. As such, the question of renormalizability is not really an issue. In [5], the method of [3, 4] was extended to two loop order. However, a kind of unitary gauge was employed, which multiplicative renormalizability is not guaranteed.

Some gauges used in the literature are nonlocal [3, 4]. Evidently, it would also remain to be seen whether the nonappearance of divergences at one loop remains the case at higher orders due to the nonlocality of the gauge, preventing a straightforward analysis of the renormalizability.

II. ULTRAVIOLET FINITENESS AT ONE LOOP ORDER.

Let us first briefly review the ultraviolet finiteness at one loop order. Let us begin with the one loop ghost-antighost self-energy. As already pointed out in [1], using the transversality of the gluon propagator in the Landau gauge, it can be checked that the one loop ghost self-energy is free from ultraviolet divergences. As a consequence we have that

$$Z_c = Z_{\bar{c}} = 1 + \mathcal{O}\left(\frac{g^4}{m^2}\right). \quad (8)$$

Analogously, the one loop correction to the ghost-antighost-gluon vertex is also finite. The same feature holds for the one loop Feynman diagrams contributing to the four-gluon vertex, from which it follows that

$$Z_g^2 Z_A^2 = 1 + \mathcal{O}\left(\frac{g^4}{m^2}\right). \quad (9)$$

Moreover, from the equations (7), we have

$$Z_A = 1 + \mathcal{O}\left(\frac{g^4}{m^2}\right), \quad (10)$$

so that

$$Z_g = 1 + \mathcal{O}\left(\frac{g^4}{m^2}\right). \quad (11)$$

We conclude that the theory is completely free from ultraviolet divergences at one loop.

III. ULTRAVIOLET FINITENESS AT TWO LOOP ORDER.

We comment briefly on the two loop computation of the renormalization constants using the massive action (5). It is based on the same approach for renormalizing the Curci-Ferrari model in four dimensions, [12, 13]. There QCD is fixed in the Curci-Ferrari gauge and the gluon and ghosts are given a mass via an on-shell BRST invariant dimension two operator. Therefore, this corresponds to a natural infrared regulator in a renormalizable theory. For our three dimensional case the mass plays a similar role as an infrared regulator. Hence we can apply the same vacuum bubble expansion method as [12, 13] inspired by the earlier approach of [14, 15]. One can systematically replace propagators involving an external momentum, p , with a propagator involving only internal momenta by the exact identity

$$\frac{1}{[(k-p)^2 + m^2]} = \frac{1}{(k^2 + m^2)} + \frac{(2kp - p^2)}{[(k-p)^2 + m^2](k^2 + m^2)}. \quad (12)$$

This can be repeated iteratively with the truncation criterion being determined by the particular Green function under consideration and exploiting the renormalizability of the Lagrangian. So, for instance, for a gluon 2-point function one expands to $\mathcal{O}(p^2)$ since terms of higher order in the external momenta will be ultraviolet finite by renormalizability and hence can be ignored for the purposes of determining the renormalization constants. The integrals which are retained involve vacuum bubbles which are reduced to Lorentz scalar integrals after an elementary reduction of the tensor structure in the numerator of the integrand. All that remains is the evaluation of these three dimensional integrals at two loops. This is achieved by the results summarized in [16] where three dimensional massive vacuum bubbles have been evaluated to *three* loops. More specifically we note that all the (non-trivial) vacuum bubbles can be related to the integral

$$\int \frac{d^d k d^d \ell}{(k^2 + m_1^2)(\ell^2 + m_2^2)[(k-\ell)^2 + m_3^2]} = \frac{1}{(4\pi)^2} \left[\frac{1}{4\varepsilon} + \frac{1}{2} + \ln \left[\frac{\mu}{(m_1 + m_2 + m_3)} \right] \right] + \mathcal{O}(\varepsilon), \quad (13)$$

where $d = 3 - 2\varepsilon$ in dimensional regularization and μ is the mass scale introduced to maintain the same dimensionality of the coupling constant in d as in three dimensions.

We have implemented the above algorithm in the symbolic manipulation language FORM, [17], and automatically renormalized all the n -point functions relevant for determining all the renormalization constants of the Lagrangian. The Feynman diagrams were generated automatically by the QGRAF package, [18], and converted to FORM input

notation. The upshot of the renormalization is that all the renormalization constants are unity up to and including two loops (in the $\overline{\text{MS}}$ scheme). At one loop this is virtually a trivial observation due to the fact that there the basic master vacuum bubble is finite

$$\int \frac{d^d k}{[k^2 + m^2]} = - \frac{m}{4\pi} \left[1 + \left[2 + 2 \ln \left(\frac{\mu}{2m} \right) \right] \varepsilon \right] + O(\varepsilon^2). \quad (14)$$

This is yet another verification of the fact that the one loop quantum corrections are finite.

At two loops the situation is less trivial since the basic integral (13) is ultraviolet divergent but when all diagrams to the Green functions involving this integral are added together then the simple pole in ε has zero residue. For instance, in the renormalization of the ghost two point function 6 of the 13 two loop Feynman diagrams are divergent but overall the resulting renormalization constant is unity.

IV. ULTRAVIOLET FINITENESS AT ALL ORDERS.

The general counterterm structure corresponding to the action (5) will look like

$$S_c = S_1 g^2 + S_2 g^4 + S_3 g^6 + S_4 \frac{g^8}{m} + \dots, \quad (15)$$

where the S_i contains the counterterms proportional to powers of $\frac{1}{\varepsilon}$. However, it is known that counterterms have to be polynomial in the fields, their derivatives and the external parameters like masses [19], when massless renormalization schemes like $\overline{\text{MS}}$ are employed. Therefore, all coefficients S_i , $i > 3$, should be zero. We can also forget about the three loop term $\propto g^6$, as this is already of dimension three, so it is merely a potentially infinite constant, but not affecting any Green function. We come to the conclusion that the only counterterms that can occur, must emerge at one or two loop order. Having explicitly checked that there are no such counterterms, we are able to conclude that the theory (5) is completely free of any ultraviolet divergence. As a consequence, all Green functions can be unambiguously calculated, without the need of any renormalization scheme.

V. THE CURCI-FERRARI GAUGE.

The mass operator $A_\mu^a A_\mu^a$ in the Landau gauge can be generalized to the Curci-Ferrari gauge, in which case the mixed gluon-ghost mass operator $(\frac{1}{2} A_\mu^a A_\mu^a + \alpha \bar{c}^a c^a)$ has to be considered, where α stands for the gauge parameter. This operator also enjoys the property of being BRST invariant on-shell. The action corresponding to this nonlinear gauge fixing reads

$$S = \int d^3 x \left(-\frac{1}{4} F_{\mu\nu}^a F_{\mu\nu}^a + b^a \partial_\mu A_\mu^a + \frac{\alpha}{2} b^a b^a + \bar{c}^a \partial_\mu D_\mu^{ab} c^b - \frac{\alpha}{2} g f^{abc} b^a \bar{c}^b c^c - \frac{\alpha}{8} g^2 f^{abc} f^{cde} \bar{c}^a \bar{c}^b c^d c^e + m^2 \left(\frac{1}{2} A_\mu^a A_\mu^a + \alpha \bar{c}^a c^a \right) \right). \quad (16)$$

The Landau gauge follows from the Curci-Ferrari gauge in the limit $\alpha \rightarrow 0$.

It was proven in [1] that (16) is multiplicatively renormalizable to all orders. Due to the Ward identities, it also turned out that the renormalization factor of the mass m is not an independent quantity of the theory either [1]. Moreover, we have checked that the results of sections III and IV do also extend to the Curci-Ferrari case. In other words, the action (16) describes an ultraviolet finite theory.

VI. CONCLUSION.

Having proven the renormalizability and in addition the ultraviolet finiteness of the actions (5) and (16), the question is what happens in other gauges, which have also been supplemented with a regulating mass that is consistent with renormalizability. We expect that all the existing results from four-dimensional Yang-Mills gauge theories should translate to three dimensions. We refer to [20, 21] for the available literature on several covariant gauges and their corresponding renormalizable mass operators. Let us also mention that these gauges and operators can be connected to each other by means of renormalizable interpolating gauges. It remains an open problem if the three-dimensional version of these mass regularized theories would also be ultraviolet finite at all orders.

We have considered regulated three-dimensional gauge theories. Of course, the real challenge would be to find a mechanism behind the dynamical generation of a mass. In principle, as it concerns a gauge theory, this mass should be obtained in a gauge invariant way. The used mass operator, $\int d^3x A_\mu^2$ is apparently not gauge invariant. Even more, it is not renormalizable in any gauge. Nevertheless, it has a special feature in the Landau gauge worth mentioning. Completely analogous as in four dimensions, A_μ^2 can be related to the operator

$$A_{\min}^2 = (VT)^{-1} \min_{U \in SU(N)} \int d^3x (A_\mu^U)^2, \quad (17)$$

which is gauge invariant due to the minimization along the gauge orbit. $A_{\min}^2$ reduces to A_μ^2 in the Landau gauge [26]. We refer to [20] for a list of references.

We also notice that the mass m should be fixed. This should be done by some kind of consistent gap equation, as m is not supposed to be a free parameter of the theory, see also [3, 4, 22]. This is however beyond the scope of the present paper.

Nevertheless, it would be interesting to pursue the search for a gauge invariant mass insertion. Inspired by [3, 4], we have started the investigation of the gauge invariant mass operator $\int d^4x F_{\mu\nu}^a \left[(D^2)^{-1} \right]^{ab} F_{\mu\nu}^b$ in four dimensions. In four dimensions, it is possible to add this operator to the action as a mass term and to localize it in a way consistent with the gauge invariance. Starting from this, a local, non-Abelian gauge invariant action with a mass was constructed [23, 24]. The renormalizability has been proven to all orders in the linear covariant gauges. This study could be extended to three dimensions, which might allow for a gauge invariant discussion of a potential dynamical, parity preserving, mass generation. An even more interesting question would be if this action would also give rise to an ultraviolet finite theory.

Acknowledgments.

The Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq-Brazil), the Faperj, Fundação de Amparo à Pesquisa do Estado do Rio de Janeiro, the SR2-UERJ and the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) are gratefully acknowledged for financial support. D. Dudal acknowledges the hospitality at the UERJ where this work was completed.

-
- [1] D. Dudal, J. A. Gracey, V. E. R. Lemes, R. F. Sobreiro, S. P. Sorella and H. Verschelde, *Annals Phys.* **317** (2005) 203.
 - [2] R. Jackiw and S. Templeton, *Phys. Rev. D* **23** (1981) 2291.
 - [3] R. Jackiw and S. Y. Pi, *Phys. Lett. B* **403** (1997) 297.
 - [4] R. Jackiw and S. Y. Pi, *Phys. Lett. B* **368** (1996) 131.
 - [5] F. Eberlein, *Phys. Lett. B* **439** (1998) 130.
 - [6] J. M. Cornwall, *Phys. Rev. D* **57** (1998) 3694.
 - [7] D. Karabali and V. P. Nair, *Nucl. Phys. B* **464** (1996) 135.
 - [8] D. Karabali and V. P. Nair, *Phys. Lett. B* **379** (1996) 141.
 - [9] O. Piguet and S. P. Sorella, *Lect. Notes Phys.* **M28** (1995) 1.
 - [10] K. I. Kondo, *Phys. Lett. B* **514** (2001) 335.
 - [11] D. Dudal, H. Verschelde and S. P. Sorella, *Phys. Lett. B* **555** (2003) 126.
 - [12] J. A. Gracey, *Phys. Lett. B* **525** (2002) 89.
 - [13] R. E. Browne and J. A. Gracey, *Phys. Lett. B* **540** (2002) 68.
 - [14] M. Misiak and M. Munz, *Phys. Lett. B* **344** (1995) 308.
 - [15] K. G. Chetyrkin, M. Misiak and M. Munz, *Nucl. Phys. B* **518** (1998) 473.
 - [16] A. K. Rajantie, *Nucl. Phys. B* **480** (1996) 729 [Erratum-ibid. B **513** (1998)].
 - [17] J. A. M. Vermaseren, *math-ph/0010025*.
 - [18] P. Nogueira, *J. Comput. Phys.* **105** (1993) 279.
 - [19] J. C. Collins, *Nucl. Phys. B* **80** (1974) 341.
 - [20] D. Dudal, M. A. L. Capri, J. A. Gracey, V. E. R. Lemes, R. F. Sobreiro, S. P. Sorella and H. Verschelde, *hep-th/0611114*.
 - [21] V. E. R. Lemes, R. F. Sobreiro and S. P. Sorella, *hep-th/0611231*.
 - [22] S. P. Sorella, *Annals Phys.* **321** (2006) 1747.
 - [23] M. A. L. Capri, D. Dudal, J. A. Gracey, V. E. R. Lemes, R. F. Sobreiro, S. P. Sorella and H. Verschelde, *Phys. Rev. D* **72** (2005) 105016.
 - [24] M. A. L. Capri, D. Dudal, J. A. Gracey, V. E. R. Lemes, R. F. Sobreiro, S. P. Sorella and H. Verschelde, *Phys. Rev. D* **74** (2006) 045008.

- [25] The exact form of the invoked mass term is also not fixed. Several possibilities exist, see [3, 4].
- [26] Finding the global minimum is however a delicate problem, due to the existence of gauge (Gribov) copies.